



Hearing Loss:

# The **#1** Most Modifiable Risk Factor of Dementia



By Dr. Keith Darrow, PhD  
Harvard and MIT Trained Neuroscientist

**Hearing  Brain**  
Centers of America



INTRODUCTORY LETTER FROM DR. DARROW

“Everyone knows someone who is a cancer survivor, but no one knows an Alzheimer’s survivor.” – Dr. Bredeesen



Dr. Keith Darrow, PhD, CCC-A

Thank you for taking the time to read this updated report detailing the links of Hearing Loss and Dementia. There are always new reports coming out helping us to better understand the science of Dementia and ways to help us avoid the devastating fate of a Dementia diagnosis. Perhaps the most impressive report I recently read was from a European Dementia commission titled: Dementia Prevention, Intervention, and Care published in the Lancet Journal.

Let’s start by reviewing some of the facts about Dementia. Nearly every 3-4 seconds another individual is diagnosed with Dementia. The average annual cost to care for a loved one with Dementia is approximately ~\$57,000 per year. Nearly 50% of all cases of Dementia are Alzheimer’s related. And, there is not a SINGLE drug available on the market approved to treat Alzheimer’s.

But, not all news is bad news. In fact, the Lancet report highlighted that nearly 35% of all Dementia cases are considered preventable. Phew! The report even laid out the most important modifiable lifestyle factors that can help each of us prevent Dementia.

And (not surprisingly), the #1, single most modifiable factor for preventing Dementia is the treatment of hearing loss. Yes, reducing obesity, diabetes, and/or cardiovascular disease are important. It is also important to increase social activity, enhance our education and supplement our nutrition. These can all play a part in helping us to prevent Dementia; but none, even when combined, are nearly as effective as the treatment of hearing loss. I think Dr. Doraiswamy, a Neuropsychologist from Duke University said it best:

“The benefits of correcting hearing loss on cognition are twice as large as the benefits from any cognitive-enhancing drugs now on the market. It should be the first thing we focus on.”

I believe you will find the information in this report helpful when wanting to learn more about your cognitive health and when it comes to choosing the right treatment plan for you. I encourage you to take advantage of our courteous offer to provide a hearing evaluation and cognitive assessment for all new patients at no-charge. Feel free to call the office to schedule your appointment.

Sincerely,

*Dr. Keith Darrow*

**Dr. Keith Darrow, Ph.D., CCC-A**  
**Harvard Medical and M.I.T. Trained Neuroscientist**  
**Certified Dementia Practitioner**

Amazon #1 New Release & Best-Selling Author  
Expert in Speech and Hearing Bioscience and Technology  
Medical Director at Hearing and Brain Centers  
Spokesman for Excellence in Audiology  
Professor at Worcester State University  
Nationally Recognized Speaker, Trainer, and Researcher  
Research Associate at Massachusetts Eye and Ear Infirmiry  
His publications and research cited over 1000 times








Figure 1: Summary of Data from Lin et al., 2011  
Johns Hopkins Medical Center

says. “Even if people feel as if they are not affected, we’re showing that it may well be a more serious problem.”

Three risk factors associated with hearing loss and Dementia include Social Isolation, Cerebral Atrophy and Cognitive Overload.

### 1. Social Isolation

Social Isolation – the impact of reduced social and physical activity. Withdrawal from social situations is common in individuals with hearing loss. Many studies cite feelings of embarrassment, fear of making mistakes in conversations, and feeling like you are not part of the conversation as the common rationale for individuals with hearing impairment to separate themselves from family, friends and community. This retreat from social activity has even been found in individuals with a mild degree of hearing loss. In addition, individuals with hearing loss are less likely to engage in physical activity. Both increased social isolation and reduced physical activity are strong risk factors for the development of Dementia.

### Active Aging: How to Reduce Social Isolation

Active Aging – the process of optimizing opportunities for better health, continuing development of knowledge, and increased security in order to maximize quality of life as you age. The word ‘active’ is used to describe a person’s involvement with social, physical, economic, spiritual and civic affairs. We all share the same goal to maintain autonomy and independence as we age, and thus we must rely on preserving the tenants of interdependence (socialization and reliance on family and loved ones) and intergenerational solidarity (maintaining companionship with age-matched peers) to ensure active aging.

Both Social Isolation and Depression are major risk factors for the development of Dementia, and both increase as we age. Being a lifelong learner and staying active is important to maintain a healthy, active brain and can also reduce your risk of cognitive decline and dementia. Some studies have shown that social activities, larger social networks and a history of social



“Blindness separates us from things, but deafness separates us from people.

- Helen Keller

contact are associated with better cognitive function and reduced risk for cognitive decline.

## 2. Cerebral Atrophy (aka Brain Shrinkage)

The association of a shrinking brain, resulting from the loss of neurons, with Dementia has been long documented. Even people with MCI (Mild Cognitive Impairment) show signs of cerebral atrophy. In recent years, scientific studies using advanced brain imaging techniques (including fMRI - Functional Magnetic Reso-


nance Imaging) have demonstrated that hearing impairment is associated with accelerated brain atrophy in both the overall brain as well as even more advanced reductions in volume associated with the memory, hearing, speech and language portions of the brain.

Individuals with Hearing Loss can experience significant cerebral atrophy. The most significant reduction in cerebral volume occurs in areas involved in:

- Memory
- Hearing
- Speech
- Language

“It (hearing loss) not only increases age-related memory loss, it increases the incidents of Alzheimer’s disease so if you can prevent the onset of Alzheimer’s disease or delay it with good hearing devices that’s a major public health advance.

- Quote by Erik Kandel, Recipient of the Nobel Prize in Physiology or Medicine



**Brain With Hearing Loss**

**Brain With Normal Hearing**

### **3. Cognitive Overload (i.e. Working Your Brain Too Hard To Hear)**

Hearing loss is not normal and neither is the excess strain that it puts on your brain. While hearing loss may be more common as we age, it is critical that hearing loss be treated. With hearing loss, the brain is constantly on 'over- load' trying to fill in the missing pieces and follow the conversation. Increased cognitive load is considered a risk factor for developing Dementia. Cognitive load, as measured by pup- illometry, is a measurement of how hard your brain is working to follow a conversation. Recent research has found that individuals who treat their hearing loss do not work as hard to listen (i.e. have a reduced cognitive load) and have as much as a 20% increase in memory recall when following a conversation.

### **NeuroTechnology™:**

#### **Reduce Your Risk of Developing Dementia**

**Improvements in Cognitive Function:** In a recent study, Dr. Jamie Desjardins, PhD (University of Texas at El Paso) demonstrated that today's current hearing loss treatment options can improve brain function in people with hearing loss. It is known that hearing loss, if left untreated, can lead to emotional and social consequences, reduced job performance and diminished quality of life. Recently, research has shown that untreated hearing loss can also can interfere with cognitive abilities because so much mental effort is diverted toward understanding speech.

The research was aimed at measuring core cognitive functions that were beginning hearing



loss treatment with NeuroTechnology™. After only two weeks of hearing loss treatment, cognitive testing revealed a significant increase in percent scores for recalling words in working memory and selective attention tests, and the processing speed at which participants selected the correct response was faster. By the end of the study, participants had exhibited significant improvement in their cognitive function. Since

2011, multiple long-term studies have provided strong evidence that treating hearing loss may eliminate the risk of developing Dementia. Dr. Lalwani at Columbia University noted that treating hearing loss “may offer a simple, yet important, way to prevent or slow the development of dementia by keeping adults with hearing loss engaged in conversation and communication.”

#### TIPS FOR ACTIVE AGING INCLUDE:

- Share a meal with family and friends 3-5 times per week
- Commit to an aerobics / exercise regimen
- Learn a new hobby each year
- Play an instrument (or learn a new instrument)
- If you love to read.... Keep reading (try to mix up the topics!)
- If you don't read much - try to read a book every other month
- Participate in classes at your local senior center
- Volunteer at a local hospital, shelter, etc.
- Go back to school. Many local Universities offer free tuition to people over 65!

*Hearing Loss:*

# The #1 Most Modifiable Risk Factor of Dementia



# Hearing Brain

Centers of America

10585 N. Tatum Blvd. Suite D-135, Paradise  
Valley, AZ 85253

8900 E Pinnacle Peak Rd. Suite C-2, Scottsdale,  
AZ 85255

18555 N. 79th Ave, Suite A101, Glendale, AZ  
85308

12691 W. Smokey Dr. Suite 131, Surprise, Arizona  
85378

5010 E. Warner Road #114, Phoenix, AZ 85044